

NAG Library Routine Document

D01FDF

Note: before using this routine, please read the Users' Note for your implementation to check the interpretation of *bold italicised* terms and other implementation-dependent details.

1 Purpose

D01FDF calculates an approximation to a definite integral in up to 30 dimensions, using the method of Sag and Szekeres (see Sag and Szekeres (1964)). The region of integration is an n -sphere, or by built-in transformation via the unit n -cube, any product region.

2 Specification

```

SUBROUTINE D01FDF (NDIM, F, SIGMA, REGION, LIMIT, RO, U, RESULT, NCALLS,
1                IFAIL)
INTEGER          NDIM, LIMIT, NCALLS, IFAIL
double precision F, SIGMA, RO, U, RESULT
EXTERNAL        F, REGION

```

3 Description

D01FDF calculates an approximation to

$$\int_{n\text{-sphere of radius } \sigma} f(x_1, x_2, \dots, x_n) dx_1 dx_2 \cdots dx_n \quad (1)$$

or, more generally,

$$\int_{c_1}^{d_1} dx_1 \cdots \int_{c_n}^{d_n} dx_n f(x_1, \dots, x_n) \quad (2)$$

where each c_i and d_i may be functions of x_j ($j < i$).

The routine uses the method of Sag and Szekeres (1964), which exploits a property of the shifted p -point trapezoidal rule, namely, that it integrates exactly all polynomials of degree $< p$ (see Krylov (1962)). An attempt is made to induce periodicity in the integrand by making a parameterized transformation to the unit n -sphere. The Jacobian of the transformation and all its direct derivatives vanish rapidly towards the surface of the unit n -sphere, so that, except for functions which have strong singularities on the boundary, the resulting integrand will be pseudo-periodic. In addition, the variation in the integrand can be considerably reduced, causing the trapezoidal rule to perform well.

Integrals of the form (1) are transformed to the unit n -sphere by the change of variables:

$$x_i = y_i \frac{\sigma}{r} \tanh\left(\frac{ur}{1-r^2}\right)$$

where $r^2 = \sum_{i=1}^n y_i^2$ and u is an adjustable parameter.

Integrals of the form (2) are first of all transformed to the n -cube $[-1, 1]^n$ by a linear change of variables

$$x_i = ((d_i + c_i) + (d_i - c_i)y_i)/2$$

and then to the unit sphere by a further change of variables

$$y_i = \tanh\left(\frac{uz_i}{1-r}\right)$$

where $r^2 = \sum_{i=1}^n z_i^2$ and u is again an adjustable parameter.

The parameter u in these transformations determines how the transformed integrand is distributed between the origin and the surface of the unit n -sphere. A typical value of u is 1.5. For larger u , the integrand is concentrated toward the centre of the unit n -sphere, while for smaller u it is concentrated toward the perimeter.

In performing the integration over the unit n -sphere by the trapezoidal rule, a displaced equidistant grid of size h is constructed. The points of the mesh lie on concentric layers of radius

$$r_i = \frac{h}{4} \sqrt{n + 8(i - 1)}, \quad i = 1, 2, 3, \dots$$

The routine requires you to specify an approximate maximum number of points to be used, and then computes the largest number of whole layers to be used, subject to an upper limit of 400 layers.

In practice, the rapidly-decreasing Jacobian makes it unnecessary to include the whole unit n -sphere and the integration region is limited by a user-specified cut-off radius $r_0 < 1$. The grid-spacing h is determined by r_0 and the number of layers to be used. A typical value of r_0 is 0.8.

Some experimentation may be required with the choice of r_0 (which determines how much of the unit n -sphere is included) and u (which determines how the transformed integrand is distributed between the origin and surface of the unit n -sphere), to obtain best results for particular families of integrals. This matter is discussed further in Section 8.

4 References

Krylov V I (1962) *Approximate Calculation of Integrals* (trans A H Stroud) Macmillan

Sag T W and Szekeres G (1964) Numerical evaluation of high-dimensional integrals *Math. Comput.* **18** 245–253

5 Parameters

- 1: NDIM – INTEGER *Input*
On entry: n , the number of dimensions of the integral.
Constraint: $1 \leq \text{NDIM} \leq 30$.
- 2: F – **double precision** FUNCTION, supplied by the user. *External Procedure*
 F must return the value of the integrand f at a given point.

The specification of F is:

```
double precision FUNCTION F(NDIM, X)
  INTEGER                NDIM
double precision      X(NDIM)
```

- 1: NDIM – INTEGER *Input*
On entry: n , the number of dimensions of the integral.
- 2: X(NDIM) – **double precision** array *Input*
On entry: the co-ordinates of the point at which the integrand f must be evaluated.

F must be declared as EXTERNAL in the (sub)program from which D01FDF is called. Parameters denoted as *Input* must **not** be changed by this procedure.

- 3: SIGMA – **double precision** *Input*
On entry: indicates the region of integration.
 SIGMA \geq 0.0
 The integration is carried out over the n -sphere of radius SIGMA, centred at the origin.
 SIGMA < 0.0
 The integration is carried out over the product region described by REGION.
- 4: REGION – SUBROUTINE, supplied by the NAG Library or the user. *External Procedure*
 If SIGMA < 0.0, REGION must evaluate the limits of integration in any dimension.

The specification of REGION is:

```
SUBROUTINE REGION(NDIM, X, J, C, D)
  INTEGER          NDIM, J
  double precision X(NDIM), C, D
```

- | | | |
|----|---|---------------|
| 1: | NDIM – INTEGER | <i>Input</i> |
| | <i>On entry:</i> n , the number of dimensions of the integral. | |
| 2: | X(NDIM) – double precision array | <i>Input</i> |
| | <i>On entry:</i> X(1), ..., X($j - 1$) contain the current values of the first ($j - 1$) variables, which may be used if necessary in calculating c_j and d_j . | |
| 3: | J – INTEGER | <i>Input</i> |
| | <i>On entry:</i> the index j for which the limits of the range of integration are required. | |
| 4: | C – double precision | <i>Output</i> |
| | <i>On exit:</i> the lower limit c_j of the range of x_j . | |
| 5: | D – double precision | <i>Output</i> |
| | <i>On exit:</i> the upper limit d_j of the range of x_j . | |

REGION must be declared as EXTERNAL in the (sub)program from which D01FDF is called. Parameters denoted as *Input* must **not** be changed by this procedure.

If SIGMA \geq 0.0, REGION is not called by D01FDF, but a dummy routine must be supplied (D01FDV may be used).

- 5: LIMIT – INTEGER *Input*
On entry: the approximate maximum number of integrand evaluations to be used.
Constraint: LIMIT \geq 100.
- 6: R0 – **double precision** *Input*
On entry: the cut-off radius on the unit n -sphere, which may be regarded as an adjustable parameter of the method.
Suggested value: a typical value is R0 = 0.8. (See also Section 8.)
Constraint: 0.0 < R0 < 1.0.
- 7: U – **double precision** *Input*
On entry: must specify an adjustable parameter of the transformation to the unit n -sphere.

Suggested value: a typical value is $U = 1.5$. (See also Section 8.)

Constraint: $U > 0.0$.

- 8: RESULT – *double precision* *Output*
On exit: the approximation to the integral I .
- 9: NCALLS – INTEGER *Output*
On exit: the actual number of integrand evaluations used. (See also Section 8.)
- 10: IFAIL – INTEGER *Input/Output*
On entry: IFAIL must be set to 0, -1 or 1. If you are unfamiliar with this parameter you should refer to Section 3.3 in the Essential Introduction for details.
On exit: IFAIL = 0 unless the routine detects an error (see Section 6).

For environments where it might be inappropriate to halt program execution when an error is detected, the value -1 or 1 is recommended. If the output of error messages is undesirable, then the value 1 is recommended. Otherwise, if you are not familiar with this parameter, the recommended value is 0. **When the value -1 or 1 is used it is essential to test the value of IFAIL on exit.**

6 Error Indicators and Warnings

If on entry IFAIL = 0 or -1 , explanatory error messages are output on the current error message unit (as defined by X04AAF).

Errors or warnings detected by the routine:

IFAIL = 1

On entry, $NDIM < 1$,
 or $NDIM > 30$.

IFAIL = 2

On entry, $LIMIT < 100$.

IFAIL = 3

On entry, $R0 \leq 0.0$,
 or $R0 \geq 1.0$.

IFAIL = 4

On entry, $U \leq 0.0$.

7 Accuracy

No error estimate is returned, but results may be verified by repeating with an increased value of LIMIT (provided that this causes an increase in the returned value of NCALLS).

8 Further Comments

The time taken by D01FDF will be approximately proportional to the returned value of NCALLS, which, except in the circumstances outlined in (b) below, will be close to the given value of LIMIT.

(a) Choice of $R0$ and U

If the chosen combination of r_0 and u is too large in relation to the machine accuracy it is possible that some of the points generated in the original region of integration may transform into points in the

unit n -sphere which lie too close to the boundary surface to be distinguished from it to machine accuracy (despite the fact that $r_0 < 1$). To be specific, the combination of r_0 and u is too large if

$$\frac{ur_0}{1-r_0^2} > 0.3465(t-1), \quad \text{if SIGMA} \geq 0.0,$$

or

$$\frac{ur_0}{1-r_0} > 0.3465(t-1), \quad \text{if SIGMA} < 0.0,$$

where t is the number of bits in the mantissa of a *double precision* number.

The contribution of such points to the integral is neglected. This may be justified by appeal to the fact that the Jacobian of the transformation rapidly approaches zero towards the surface. Neglect of these points avoids the occurrence of overflow with integrands which are infinite on the boundary.

(b) Values of LIMIT and NCALLS

LIMIT is an approximate upper limit to the number of integrand evaluations, and may not be chosen less than 100. There are two circumstances when the returned value of NCALLS (the actual number of evaluations used) may be significantly less than LIMIT.

Firstly, as explained in (a), an unsuitably large combination of R0 and U may result in some of the points being unusable. Such points are not included in the returned value of NCALLS.

Secondly, no more than 400 layers will ever be used, no matter how high LIMIT is set. This places an effective upper limit on NCALLS as follows:

| | |
|-----------|---------|
| $n = 1 :$ | 56 |
| $n = 2 :$ | 1252 |
| $n = 3 :$ | 23690 |
| $n = 4 :$ | 394528 |
| $n = 5 :$ | 5956906 |

9 Example

This example calculates the integral

$$\int \int \int_s \frac{dx_1 dx_2 dx_3}{\sqrt{\sigma^2 - r^2}} = 22.2066$$

where s is the 3-sphere of radius σ , $r^2 = x_1^2 + x_2^2 + x_3^2$ and $\sigma = 1.5$. Both sphere-to-sphere and general product region transformations are used. For the former, we use $r_0 = 0.9$ and $u = 1.5$; for the latter, $r_0 = 0.8$ and $u = 1.5$.

9.1 Program Text

```
*      D01FDF Example Program Text
*      Mark 14 Revised. NAG Copyright 1989.
*      .. Parameters ..
      INTEGER          NOUT
      PARAMETER       (NOUT=6)
*      .. Local Scalars ..
      DOUBLE PRECISION R0, RESULT, SIGMA, U
      INTEGER          IFAIL, LIMIT, NCALLS, NDIM
*      .. External Functions ..
      DOUBLE PRECISION F
      EXTERNAL        F
*      .. External Subroutines ..
      EXTERNAL        D01FDF, D01FDV, REGION
*      .. Executable Statements ..
      WRITE (NOUT,*) 'D01FDF Example Program Results'
      NDIM = 3
      LIMIT = 8000
      U = 1.5D0
      WRITE (NOUT,*)
```

```

WRITE (NOUT,*) 'Sphere-to-sphere transformation'
SIGMA = 1.5D0
RO = 0.9D0
IFAIL = 1
*
CALL D01FDF(NDIM,F,SIGMA,D01FDV,LIMIT,RO,U,RESULT,NCALLS,IFAIL)
*
IF (IFAIL.EQ.0) THEN
  WRITE (NOUT,*)
  WRITE (NOUT,99999) 'Estimated value of the integral =', RESULT
  WRITE (NOUT,99998) 'Number of integrand evaluations =', NCALLS
  WRITE (NOUT,*)
  WRITE (NOUT,*) 'Product region transformation'
  SIGMA = -1.0D0
  RO = 0.8D0
  IFAIL = 0
*
  CALL D01FDF(NDIM,F,SIGMA,REGION,LIMIT,RO,U,RESULT,NCALLS,IFAIL)
*
  WRITE (NOUT,*)
  WRITE (NOUT,99999) 'Estimated value of the integral =', RESULT
  WRITE (NOUT,99998) 'Number of integrand evaluations =', NCALLS
ELSE
  WRITE (NOUT,*)
  WRITE (NOUT,99997) ' ** D01FDF returned with IFAIL = ', IFAIL
END IF
*
99999 FORMAT (1X,A,F9.3)
99998 FORMAT (1X,A,I4)
99997 FORMAT (1X,A,I5)
END
*
DOUBLE PRECISION FUNCTION F(NDIM,X)
*
.. Scalar Arguments ..
INTEGER      NDIM
*
.. Array Arguments ..
DOUBLE PRECISION X(NDIM)
*
.. Local Scalars ..
INTEGER      I
*
.. Intrinsic Functions ..
INTRINSIC    ABS, SQRT
*
.. Executable Statements ..
F = 2.25D0
DO 20 I = 1, NDIM
  F = F - X(I)*X(I)
20 CONTINUE
F = 1.0D0/SQRT(ABS(F))
RETURN
END
*
SUBROUTINE REGION(NDIM,X,J,C,D)
*
.. Scalar Arguments ..
DOUBLE PRECISION C, D
INTEGER      J, NDIM
*
.. Array Arguments ..
DOUBLE PRECISION X(NDIM)
*
.. Local Scalars ..
DOUBLE PRECISION SUM
INTEGER      I, J1
*
.. Intrinsic Functions ..
INTRINSIC    ABS, SQRT
*
.. Executable Statements ..
C = -1.5D0
D = 1.5D0
IF (J.GT.1) THEN
  SUM = 2.25D0
  J1 = J - 1
  DO 20 I = 1, J1
    SUM = SUM - X(I)*X(I)
20 CONTINUE

```

```
D = SQRT(ABS(SUM))  
C = -D  
END IF  
RETURN  
END
```

9.2 Program Data

None.

9.3 Program Results

D01FDF Example Program Results

Sphere-to-sphere transformation

Estimated value of the integral = 22.168
Number of integrand evaluations =8026

Product region transformation

Estimated value of the integral = 22.137
Number of integrand evaluations =8026
